

Coelogyne pangasinanense 'Arnie' CBR/AOS 0 pts. 20235780

One hundred fifty-nine broadly stellate flowers nicely displayed on eight long, strongly arching, up to 20.0-cm inflorescences; plant very clean, well grown, in New Zealand sphagnum moss in a 15.0-cm hexagonal black plastic basket; leaves elliptic-oblong, light green, heavily veined, acute, petiolate 6.0 cm, originates from apex of the pseudobulb; pseudobulb spindle-shaped, 4.0 cm long by 1.6 cm diameter, covered by thin, dried sheaths; Inflorescences racemic, emerging from newly developing growths, originating from the apex of the new pseudobulb; sepals and petals white transition to light yellow-green, darker distally, sepals lanceolate, petals oblanceolate; lip cream, overlaid green-yellow, side lobes and margin cream, midlobe broadly ovate; column white, blushed broadly along midvein on dorsal surface, anther cap cream; substance firm; texture matte; country of origin is the Philippines; pending SITF confirmation; recognized for pleasant sprays of yellow-green, stellate flowers on each of multiple arching inflorescences.

Additional Reference information:

Flower Size 3/8" [9mm]

Found only in the Philippines on rocks along roadsides in full sun at elevations of 600 to 900 meters as a *miniature sized, warm growing lithophyte with erect, slender, semi-spindle-shaped, yellowish in maturity pseudobulbs subtended by elongate soon fibrous sheaths and carrying elliptic-oblong, heavily veined, petiolate, terete below the middle leaf that envelops the inflorescence when first arising [synanthous] as a slender erect, relatively stout, terete to somewhat compressed, subglabrous, and becomes 10" [25 cm] long, racemose inflorescence with 12 to many distichously alternating, fragrant flowers in a nodding to pendant rachis occurring in the late spring.*